arXiv:1703.07010v1 [math.NT] 21 Mar 2017

ON FROBENIUS AND FIBERS OF ARITHMETIC JET SPACES

JAMES BORGER AND ARNAB SAHA

Abstract. In this article, given a scheme X we show the existence of canonical lifts of Frobenius maps in an inverse system of schemes obtained from the fiber product of the canonical prolongation sequence of arithmetic jet spaces J^*X and a prolongation sequence S^* over the scheme X. As a consequence, for any smooth group scheme E, if N^n denote the kernel of the canonical projection map of the n-th jet space $J^nE \to E$, then the inverse system $\{N^n\}_n$ is a prolongation sequence.

1. Introduction

The purpose of this short note is to make an observation which is a generalisation of a result shown in [3] for Drinfeld modules. Let B be a Dedekind domain and fix a maximal ideal $\mathfrak{p} \in \operatorname{Spec} B$ with $k := B/\mathfrak{p}$ a finite field and let $q = |k|$. Let R be the $\mathfrak{p}$-adic completion of B. Denote by $\mathfrak{m}$ the maximal ideal of the complete, local ring R and $\iota : B \hookrightarrow R$ the natural inclusion. Then let $\pi \in B$ be such that $\iota(\pi)$ generates the maximal ideal $\mathfrak{m}$ in R. Since ι is an injection, by abuse of notation, we will consider π as an element of B as well. Then $k \simeq R/(\pi)$. Do note here that the identity map on R lifts the q-power Frobenius on $R/(\pi)$. Let $V = \operatorname{Spec} R$.

To motivate the main result in this article, let E be a group scheme over V. Then by [2, 4], one can consider, for all integer $n \geq 0$, the n-th jet space J^nE (Here by J^nE, we understand the algebraic jet space in [2]. Buium's jet spaces are p-adic formal schemes obtained by taking formal completions of J^nE). Due to functorial reasons, J^nE is also a group scheme for all n. There are canonical maps, the projection map $u : J^nE \to J^{n-1}E$ and the lift of Frobenius with respect to u denoted $\phi : J^nE \to J^{n-1}E$ for all $n \geq 1$. The map ϕ is associated to a π-derivation δ on the structure sheaves which will be defined in section 2 in detail. Such a system of schemes is called a *prolongation sequence*. Let J^*E denote the prolongation sequence of jet spaces. Then for each n, we have the following short exact sequence of group schemes

$$0 \longrightarrow N^n \longrightarrow J^nE \xrightarrow{u} E \longrightarrow 0 \tag{1.1}$$

where N^n is the kernel of u. Then it is easy to see that the projection map $u : J^nE \to J^{n-1}E$ induces a $u : N^n \to N^{n-1}$ by restriction. Let N^* denote the inverse system of such schemes. However, the δ or ϕ does not restrict to N^* because that would imply that E has a lift of Frobenius which is false in general.

Let X be a scheme over V. For any prolongation sequences T^* and S^* over V with morphisms $T^0 \to X$ and $S^0 \to X$, let us define $T^* \times_X S^* := \{T^n \times_X S^n\}_{n=0}^{\infty}$. Note that $T^* \times_X S^*$ is not apriori a prolongation sequence but is an inverse system

with the projection maps induced from the projection maps from T^* and S^*. For any inverse system of schemes U^* with maps $u : U^n \to U^{n-1}$ for all $n \geq 1$, we will say U^* admits a lift of Frobenius if for all n, there are maps $\phi : U^n \to U^{n-1}$ which are lifts of Frobenius with respect to the projection maps u. Our main result is

Theorem 1.1. *If X is a scheme over V, $J^*X \times_X S^*$ admits a canonical lift of Frobenius.*

We call this canonical lift of Frobenius as *Lateral Frobenius* and is denoted by $\mathfrak{f} : J^nX \times_X S^n \to J^{n-1}X \times_X S^{n-1}$ for all $n \geq 1$. Then if E is a smooth group scheme, then J^nE are also smooth and hence flat. Therefore a lift of Frobenius is equivalent to a π-derivation in this case. It is now easy to see that by choosing $X = E$ and S^* to be the constant prolongation sequence given by $S^n := V$ for all n, with the given map $S^0 \to E$ as the identity section of the group scheme E, it follows that N^n admits a lift of Frobenius and thus is a prolongation sequence.

2. Prelimineries

2.1. **Witt vectors.** Here we recall some basic facts about Witt vectors and arithmetic jet spaces. Witt vectors over a general Dedekind domain with finite residue fields was developed in [1]. For the sake of our article, we will briefly review the general construction. Let B be a Dedekind domain and fix a maximal ideal $\mathfrak{p} \in \operatorname{Spec} B$ with $k := B/\mathfrak{p}$ a finite field and let $q = |k|$. Let R be the $\mathfrak{p}$-adic completion of B. Denote by $\mathfrak{m}$ the maximal ideal of the complete, local ring R and $\iota : B \hookrightarrow R$ the natural inclusion. Then let $\pi \in B$ be such that $\iota(\pi)$ generates the maximal ideal $\mathfrak{m}$ in R. Since ι is an injection, by abuse of notation, we will consider π as an element of B as well. Then $k \simeq R/(\pi)$.

Do note here that the identity map on R lifts the q-power Frobenius on $R/(\pi)$. We will now review the theory of π-typical Witt vectors over R with maximal ideal $\mathfrak{m}$. All the rings in this section are R-algebras.

Let C be an A-algebra with structure map $u : A \to C$. In this paper, any ring homomorphism $\psi : A \to C$ will be called the *lift of Frobenius* if it satisfies the following:

(1) The reduction mod π of ψ is the q-power Frobenius, that is, $\psi(x) \equiv u(x)^q \mod \pi C$.

(2) The restriction of ψ to R is identity.

Let C be an A-algebra with structure map $u : A \to C$. A π-derivation δ from A to C means a set theoretic map satisfying the following for all $x, y \in A$

$$\begin{aligned} \delta(x+y) &= \delta(x) + \delta(y) + C_\pi(u(x), u(y)) \\ \delta(xy) &= u(x)^q\delta(y) + u(y)^q\delta(x) + \pi\delta(x)\delta(y) \end{aligned}$$

such that δ when restricted to R is $\delta(r) = (r - r^q)/\pi$ for all $r \in R$ and

$$C_\pi(X,Y) = \begin{cases} 0, & \text{if } R \text{ is positive characteristic} \\ \frac{X^q + Y^q - (X+Y)^q}{\pi}, & \text{otherwise} \end{cases}$$

It follows that the map $\phi : A \to C$ defined as

$$\phi(x) := u(x)^q + \pi\delta(x)$$

is an R-algebra homomorphism and is a lift of the Frobenius. Considering this operator δ leads to Buium's theory of arithmetic jet spaces [4, 5, 6].

Note that this definition depends on the choice of uniformizer π, but in a transparent way: if π' is another uniformizer, then $\delta(x)\pi/\pi'$ is a π'-derivation, and this correspondence induces a bijection between π-derivations and π'-derivations.

Given an R-algebra A, the ring of π-typical Witt vectors $W(A)$ can be defined as the unique R-algebra $W(A)$ with a π-derivtion δ on $W(A)$ such that, given any R-algebra C with a π-derivation δ on it and an R-algebra map $f : C \to A$, there exists an unique R-algebra homomorphism $g : C \to W(A)$ satisfying-

$$\begin{array}{ccc} W(B) & & \\ \downarrow & \overset{g}{\nwarrow} & \\ A & \xleftarrow{f} & C \end{array}$$

and g satisfies $g \circ \delta = \delta \circ g$. In [1] (following the approach of [7] to the usual p-typical Witt vectors), the existence of such a $W(A)$ is shown and that it is also obtained from the classical definition of Witt vectors using ghost vectors.

2.2. **Prolongation sequences and Jet spaces.** Let $V = \operatorname{Spec} R$ and X and Y be schemes over V. We say a pair (u, δ) is a *prolongation*, and write $Y \overset{(u,\delta)}{\to} X$, if $u : Y \to X$ is a map of schemes over V and $\delta : \mathcal{O}_X \to u_*\mathcal{O}_Y$ is a π-derivation satisfying

$$\begin{array}{ccc} B & \longrightarrow & u_*\mathcal{O}_Y \\ {\scriptstyle\delta}\uparrow & & \uparrow{\scriptstyle\delta} \\ B & \longrightarrow & \mathcal{O}_X \end{array}$$

Following [5], a *prolongation sequence* is a sequence of prolongations

$$V \xleftarrow{(u,\delta)} T^0 \xleftarrow{(u,\delta)} T^1 \xleftarrow{(u,\delta)} \cdots,$$

where each T^n is a scheme over V. We will often use the notation T^* or $\{T_n\}_{n\geq 0}$. Note that if the T^ns are flat over V then a π-derivation δ is equivalent to a lift of Frobenius ϕ as defined above.

Prolongation sequences form a category $\mathcal{C}_{V^*}$, where a morphism $f : T^* \to U^*$ is a family of morphisms $f^n : T^n \to U^n$ commuting with both the u and δ, in the evident sense. This category has a final object V^* given by $V^n = \operatorname{Spec} R$ for all n, where each u is the identity and each δ is the given π-derivation on R.

For any V-scheme Y, for all $n \geq 0$ we define the n-th jet space J^nX (relative to V) as

$$J^nX(Y) := \operatorname{Hom}_\delta(W_n^*(Y), X)$$

where $W_n^*(Y)$ is defined as in [2]. We will not define $\operatorname{Hom}_\delta(W_n^*(Y), X)$ in full generality here. Instead, we will define $\operatorname{Hom}_\delta(W_n^*(Y), X)$ in the affine case which is obviously simpler but will offer an intuitive understanding of the definition. Let our

schemes be affine and $X = \operatorname{Spec} A$ and $Y = \operatorname{Spec} C$. Then $W_n^*(Y) = \operatorname{Spec} W_n(C)$ and $\operatorname{Hom}_\delta(W_n^* Y, X) = \operatorname{Hom}_\delta(A, W_n(C))$, where $W_n(C)$ is the ring of truncated Witt vectors of length $n+1$ and $f \in \operatorname{Hom}_\delta(A, W_n(C))$ is a ring homomorphism that satisfies the following

$$\begin{array}{ccc} A & \xrightarrow{f} & W_n(C) \\ \uparrow & & \uparrow \\ R & \xrightarrow[\exp_\delta]{} & W_n(R) \end{array}$$

where $R \xrightarrow{\exp_\delta} W_n(R)$ is the universal map from the definition of Witt vectors.

Then $J^*X := \{J^n X\}_{n\geq 0}$ forms a prolongation sequence and is called the *canonical prolongation sequence*. By [5], J^*X satisfies the following universal property—for any $T^* \in \mathcal{C}_{V^*}$ and X a scheme, we have

$$\operatorname{Hom}(S^0, X) \simeq \operatorname{Hom}_{\mathcal{C}_{S^*}}(S^*, J^*X)$$

3. Construction of the Lateral Frobenius

Let all our schemes be over a base V which has a lift of Frobenius. Let $S^* = \{S^n\}_{n=0}^\infty$ be a prolongation sequence with a morphism $S^0 \to X$. Also, given a prolongation sequence, let $S^{*-1} := \{S^{n-1}\}_{n=1}^\infty$. For any prolongation sequence T^* with a morphism $T^0 \to X$, let us define $T^* \times_X S^* := \{T^n \times_X S^n\}_{n=0}^\infty$. Note that $T^* \times_X S^*$ is not apriori a prolongation sequence.

3.1. **Affine N-space case.** Let $X = \mathbb{A}^N$ where N may represent an arbitrary cardinality. For any V-scheme T, let $\prod_j^{j+m} T := \underbrace{T \times_V \cdots \times_V T}_{m+1-\text{times}}$. Then by definition of jet spaces, we have a canonical isomorphism of schemes $J^n X \simeq W_n^N$, where W_n is the truncated Witt vectors of length $n+1$. Therefore as a scheme, $J^n X$ can be canonically identified as $J^n X \simeq \prod_0^n X$. Hence $J^n X \times_X S^n \simeq (\prod_1^n X) \times_V S^n$. Let $w : J^n X \to \prod_0^n X$ be the ghost map of the product of Witt vectors and the right-hand side is referred to as the ghost components.

We define the *lateral Frobenius* $\mathfrak{f}$ as the unique morphism which makes the following diagram commutative

$$\begin{array}{ccccc} J^n X \times_X S^n & \xrightarrow{w\times \mathbb{1}} & (\prod_0^n X) \times_X S^n & \xrightarrow{\sim} & (\prod_1^n X) \times_V S^n \\ \downarrow \mathfrak{f} & & & & \downarrow \mathfrak{f} \\ J^{n-1} X \times_X S^{n-1} & \xrightarrow{w\times \mathbb{1}} & (\prod_0^{n-1} X) \times_X S^{n-1} & \xrightarrow{\sim} & (\prod_1^{n-1} X) \times_V S^{n-1} \end{array}$$

where $\mathfrak{f} : (\prod_1^n X) \times_V S^n \to (\prod_1^{n-1} X) \times_V S^{n-1}$ is given by the left-shift operator on the ghost components $\mathfrak{f}((w_1, \cdots, w_n), s) = ((w_2, ..., w_n), \phi(s))$ and completely determines the map $\mathfrak{f} : J^n X \times_X S^n \to J^{n-1} X \times_X S^{n-1}$. The map $\mathfrak{f}$ is unique and is given by $\mathfrak{f}(x_1, \cdots, x_n, s) = (z_1, \cdots, z_{n-1}, \phi(s))$ where $(x_1, \cdots, x_n) \mapsto (z_1, \cdots, z_{n-1})$ is the Frobenius map of Witt vectors and hence the map is a lift of Frobenius as well. It is clear that $\mathfrak{f}$ behaves functorially with respect to X.

Let $l : J^nX \times_X S^n \to J^nX \times_V S^n$ denote the natural map induced from the structure map $X \to V$ for all n.

Proposition 3.1. *The lateral Frobenius* $\mathfrak{f}$ *satisfies the following commutative diagram for all* $n \geq 2$

$$\begin{array}{ccc} J^nX \times_X S^n & \xrightarrow{\quad l \quad} & J^nX \times_V S^n \\ \downarrow{\scriptstyle \mathfrak{f}} & & \downarrow{\scriptstyle \phi\times\phi} \\ J^{n-1}X \times_X S^{n-1} & & J^{n-1}X \times_V S^{n-1} \\ & \searrow^{(\phi\times\phi)\circ l} & \downarrow{\scriptstyle \phi\times\phi} \\ & & J^{n-2}X \times_V S^{n-2} \end{array}$$

Proof. This follows by checking commutativity on the ghost component since w is injective $\square$

3.2. **The case of a general affine scheme.** Let $S^* = \{S^n\}_n$ be a prolongation sequence with a morphism $S^0 \xrightarrow{a} X$. We will denote this data as $S^* \xrightarrow{a} X$. Let X be an affine scheme. Then it satisfies an equaliser diagram

$$X \xrightarrow{f} Y \underset{h}{\overset{g}{\rightrightarrows}} Z$$

where $Y = \mathbb{A}^N$ and $Z = \mathbb{A}^M$, N and M are arbitrary and f is injective. Since the jet space functor J^n preserves equalisers [2] we obtain an equaliser of the corresponding jet spaces with the following commutative diagram



Lemma 3.2. *The map* $J^*X \times_Y S^* \xrightarrow{f} J^*Y \times_Y S^*$ *is injective.*

Proof. If $f(x,s) = f(x',s') \iff (f(x),s) = (f(x'),s')$ satisfying $u \circ f(x) = f \circ a(s) = u \circ f(x') = f \circ a(s')$ in Y. This implies that $x = x'$ since f is injective and $s = s'$ and we are done. $\square$

Lemma 3.3. *We have* $J^*X \times_Y S^* = J^*X \times_X S^*$

Proof. We have $(x,s) \in J^*X \times_Y S^* \iff f \circ \pi(x) = f \circ a(s)$ in $Y \iff \pi(x) = a(s)$ in X (Since f is injective) $\iff (x,s) \in J^*X \times_X S^*$ and we are done. $\square$

Combining the above two results we get an injective map $J^*X \times_X S^* \xrightarrow{f} J^*Y \times_Y S^*$.

Proposition 3.4. *The following*

$$J^*X \times_X S^* \xrightarrow{f} J^*Y \times_Y S^* \underset{h}{\overset{g}{\rightrightarrows}} J^*Z \times_Z S^*$$

is an equaliser diagram.

Proof. We need to show exactnes, that is, if for a $(y,s) \in J^*Y \times_Y S^*$ such that $g(y,s) = h(y,s)$ then we claim that $(y,s) \in J^*X \times_X S^*$. Now $g(y,s) = h(y,s) \iff (g(y),s) = (h(y),s) \iff g(y) = h(y)$ in $J^*Z \iff y \in J^*X$ (Since J^* preserves an equaliser diagram). Therefore $(y,s) \in J^*X \times_X S^*$ and we are done. □

Corollary 3.5. *If X is affine, then $J^*X \times_X S^*$ admits the lift of Frobenius induced from* $\mathfrak{f}$.

Proof. Since $Y = \mathbb{A}^N$ and $Z = \mathbb{A}^M$, we have shown that $J^*Y \times_Y S^*$ and $J^*Z \times_Z S^*$ admit the lift of Frobenius $\mathfrak{f}$. Then the result follows from the universal property of equalisers

$$\begin{array}{ccccc} J^nX \times_X S^n & \longrightarrow & J^nY \times_Y S^n & \overset{f}{\rightrightarrows} & J^nZ \times_Z S^n \\ \vdots\,\mathfrak{f} & & \downarrow\mathfrak{f} & & \downarrow\mathfrak{f} \\ J^{n-1}X \times_X S^{n-1} & \longrightarrow & J^{n-1}Y \times_Y S^{n-1} & \overset{f}{\rightrightarrows} & J^{n-1}Z \times_Z S^{n-1} \end{array}$$

□

3.3. **The case of a general scheme.** Let X be a scheme which admits a finite cover by affines X_i. Let $Y = \coprod X_i$ and $Z = Y \times_X Y$. A prolongation sequence such that $S^* \xrightarrow{a} X$, induces the map $S^* \to J^*X$ of prolongation sequences by the universal property of jet spaces. Define a new prolongation sequence $\tilde{S}^*$ given by $\tilde{S}^n := S^n \times_{J^nX} J^nY$ for all n. Define $\bar{S}^* := \tilde{S}^* \times_{S^*} \tilde{S}^*$. Then we have the following commutative diagram of coequalisers.



Proposition 3.6. *$J^*Y \times_Y \tilde{S}^* \to J^*X \times_X S^*$ is an etale surjection, that is, $J^nY \times_Y \tilde{S}^n \to J^nX \times_X S^n$ is an etale surjection for all n.*

Proof. By [2], since $Y \to X$ is etale, $J^*Y \to J^*X$ and $\tilde{S}^* \to S^*$ are etale as well. Therefore, $J^*Y \times_Y \tilde{S}^* \to J^*X \times_X S^*$ is etale.

For any V-scheme T, if $Q \to V$ is the generic point, then denote $T_Q := T \times_V Q$. If $P \to V$ is the closed point, then denote $\overline{T} := T \times_V P$. Now for all n, $(J^nY \times_Y \tilde{S}^n)_Q = (\prod_1^n Y_Q) \times \tilde{S}^n_Q$ which clearly surjects over $(J^nX \times S^n)_Q = (\prod_1^n X_Q) \times S^n_Q$. By [4], we

have $\overline{J^nY} = \overline{J^nX} \times_{\overline{X}} \overline{Y}$. Therefore, $\overline{J^nY} \times_{\overline{Y}} \overline{\tilde{S}^n} = \overline{J^nX} \times_{\overline{X}} \overline{Y} \times_{\overline{Y}} \overline{\tilde{S}^n} = \overline{J^nX} \times_{\overline{X}} \overline{\tilde{S}^n}$ which clearly surjects over $\overline{J^nX} \times_{\overline{X}} \overline{S^n}$. Hence $J^nY \times_Y \tilde{S}^n \to J^nX \times_X S^n$ is a surjection of schemes for all n and we are done. □

Proposition 3.7. $(J^*Y \times_Y \tilde{S}^*) \times_{J^*X \times_X S^*} (J^*Y \times_Y \tilde{S}^*) = (J^*Y \times_{J^*X} J^*Y) \times_{(Y \times_X Y)} (\tilde{S}^* \times_{S^*} \tilde{S}^*)$

Proof. Let $b = g \circ a$. Define a map $(J^*Y \times_Y \tilde{S}^*) \times_{J^*X \times_X S^*} (J^*Y \times_Y \tilde{S}^*) \to (J^*Y \times_{J^*X} J^*Y) \times_{(Y \times_X Y)} (\tilde{S}^* \times_{S^*} \tilde{S}^*)$ as $((y, \tilde{s}), (y', \tilde{s}')) \mapsto ((y, y'), (\tilde{s}, \tilde{s}'))$. We claim that the above map is well-defined. Consider $((y, \tilde{s}), (y', \tilde{s}')) \in (J^*Y \times_Y \tilde{S}^*) \times_{J^*X \times_X S^*} (J^*Y \times_Y \tilde{S}^*)$. Then it satisfies the following- (i) $u(y) = b(\tilde{s}), u(y') = b(\tilde{s}')$ in Y and (ii) $f(y, \tilde{s}) = f(y', \tilde{s}') \Rightarrow f(y) = f(y')$ in J^*X and $f \circ b(\tilde{s}) = f \circ b(\tilde{s}')$. But then the above two conditions implies (i) $f(y) = f(y')$ in J^*X and $f(\tilde{s}) = f(\tilde{s}')$ in S^* and (ii) $u(y) = b(\tilde{s}),\ u(y') = b(\tilde{s}')$ which implies $((y, y'), (\tilde{s}, \tilde{s}')) \in (J^*Y \times_{J^*X} J^*Y) \times_{(Y \times_X Y)} (\tilde{S}^* \times_{S^*} \tilde{S}^*)$ and hence proves the claim of well-definedness. Clearly this map has an inverse and we are done. □

Corollary 3.8.

$$J^*Z \times_Z \bar{S}^* \underset{h}{\overset{g}{\rightrightarrows}} J^*Y \times_Y \tilde{S}^* \xrightarrow{f} J^*X \times_X S^*$$

is a coequaliser diagram.

Proof. Since $\bar{S}^* = \tilde{S}^* \times_{S^*} \tilde{S}^*$, the result follows from proposition 3.6 and 3.7. □

Proof of Theorem 1.1 Since Z and Y are affine, by corollary 3.5, we know that $J^*Z \times_Z S^*$ and $J^*Y \times_Y S^*$ admit the lift of Frobenius $\mathfrak{f}$ and hence the result for $J^*X \times_X S^*$ follows from the universal property of coequalisers,

$$\begin{array}{ccccc}
J^nZ \times_Z \bar{S}^n & \underset{h}{\overset{g}{\rightrightarrows}} & J^nY \times_Y \tilde{S}^n & \xrightarrow{f} & J^nX \times_X S^n \\
\downarrow \mathfrak{f} & & \downarrow \mathfrak{f} & & \vdots \mathfrak{f} \\
J^{n-1}Z \times_Z \bar{S}^{n-1} & \underset{h}{\overset{g}{\rightrightarrows}} & J^{n-1}Y \times_Y \tilde{S}^{n-1} & \xrightarrow{f} & J^{n-1}X \times_X S^{n-1}
\end{array}$$

□

References

[1] J. Borger. The basic geometry of Witt vectors I: The affine case. *Algebra & Number Theory*, 5(2):231–285, 2011.

[2] J. Borger. The basic geometry of Witt vectors II: Spaces. *Mathematische Annalen*, 351(4):877–933, 2011.

[3] J. Borger and A. Saha. Differential Characters of Drinfeld Modules and de Rham Cohomology. *http://arxiv.org/abs/1703.05677*.

[4] A. Buium. Differential characters of abelian varieties over p-adic fields. *Inventiones mathematicae*, 122(1):309–340, 1995.

[5] A. Buium. Differential modular forms. *Journal fur die Reine und Angewandte Mathematik*, 520:95–168, 2000.

[6] A. Buium and B. Poonen. Independence of points on elliptic curves arising from special points on modular and shimura curves, II: local results. *Compositio Mathematica*, 145(03):566–602, 2009.

[7] André Joyal. δ-anneaux et vecteurs de Witt. *CR Math. Rep. Acad. Sci. Canada*, 7(3):177–182, 1985.